

Recent Improvements in Surface-Only Flux Algorithms (SOFA) (Simple Surface Flux Models)

Shashi K. Gupta², David P. Kratz¹,
Anne C. Wilber², and Victor E. Sothcott²

¹NASA Langley Research Center

²Science Systems and Applications, Inc.

Eighth CERES-II Science Team Meeting

held jointly with

Canadian Cloud-Aerosol Feedbacks and Climate Network Workshop

and

Aerosol Indirect Effects Workshop

Victoria, BC, Canada

12-16 November 2007



Climate Science Branch, NASA Langley Research Center



Background

- CERES uses several simple surface flux models for computing SW and LW surface fluxes in addition to the detailed model used by SARB. These models are:

LPSA/LPLA:
Langley Parameterized
SW/LW Algorithm

		Model A	Model B	Model C
SW	Clear	Li et al.	LPSA	--
	All-Sky	--	LPSA	--
LW	Clear	Inamdar and Ramanathan	LPLA	Zhou-Cess
	All-Sky	--	LPLA	Zhou-Cess

References:

SW A: Li et al. (1993): *J. Climate*, **6**, 1764-1772.

SW B: Darnell et al. (1992): *J Geophys. Res.*, **97**, 15741-15760.


Gupta et al. (2001): *NASA/TP-2001-211272*, 31 pp.

LW A: Inamdar and Ramanathan (1997): *Tellus*, **49B**, 216-230.

LW B: Gupta et al. (1992): *J. Appl. Meteor.*, **31**, 1361-1367.


LW C: Zhou and Cess (2001): *J. Geophys. Res.*, **106**, 12477-12488.

Zhou et al. (2007): *J. Geophys. Res.*, **112**, D15102.




Background (contd.)

- These models are fast parameterizations for TOA-to-Surface transfer algorithms; have been around and have been validated.
- The purpose of these models is to produce surface fluxes quickly, and in the process, keep a check on SARB results.
- SW and LW Models A and B used since the beginning. Good results and validation shown for all Editions of TRMM and Terra data at earlier STMs. Also for Aqua-1A and -1B data.
- LW Model C introduced recently; Tested thoroughly. Will be used starting with Edition-3 processing.
- Focus of this presentation concerns improvements made recently to all three LW models (A, B, and C), which were overestimating DLF for some footprints at the high end of DLF range.



Overestimation of DLF Over Desert Regions

Clear-Sky (Aqua-2A; July 2002 - March 2005)





Climate Science Branch, NASA Langley Research Center



Overestimation of DLF Over Desert Regions

All-Sky (Aqua-2A; July 2002 - March 2005)



Climate Science Branch, NASA Langley Research Center



Cause of Overestimation and Remedy

- Started to look for the cause of overestimation in LW Model B.
- Determine the cause, develop remedies, and apply to all models.
- Effective Emitting Temperature:
$$T_{\text{eff}} = 0.60 T_s + 0.35 T_1 + 0.05 T_2$$
 - T_s - Surface skin temperature
 - T_1 - Average temperature for Sfc. - 800 mb layer
 - T_2 - Average temperature for 800 - 680 mb layer
- Works well when T_s , T_1 , and T_2 conform to nominal lapse rates.
- When $T_s \gg T_1$ (and T_2), it results in overestimation of DLF.
- Happened mostly over dry/arid regions during hot times of the day.




Further Study of Overestimation

- Selected two sites for study:


Alice Springs, Australia - Dry/Arid

Goodwin Creek, MS (USA) - Moderate/Humid


- Performed flux computations using an offline version of the model on a 3-hourly time resolution for all months of 2000.
- Compared model-derived DLF with ground-based observations for the above sites obtained from BSRN database.



Model-Derived vs. Ground-Measured BSRN DLF (Offline LW Model B)





Significant overestimation over Alice Springs; almost none over Goodwin Creek



Climate Science Branch, NASA Langley Research Center



Model-Derived vs. Ground-Measured DLF (Offline LW Model B) Alice Springs



Severe overestimation during DJF; slight underestimation during JJA




Climate Science Branch, NASA Langley Research Center





Analysis of Alice Springs Overestimation


- Points with overestimation of $> 100 \text{ Wm}^{-2}$
32 points during the year (16 in Jan; 20 in DJF)
Mean = 120 Wm^{-2} ; Range: $100\text{-}160 \text{ Wm}^{-2}$
 T_s : Mean = 324.3 K ; Range: $302\text{-}336 \text{ K}$
 P_s : Mean = 939 mb ; Range: $935\text{-}946 \text{ mb}$
 T_{800} : Mean = 292.1 K ; Range: $284.7\text{-}297.7 \text{ K}$
 $T_s - T_{800}$: Mean = 32.2 K ; Range: $17.0\text{-}40.3 \text{ K}$
- $T_s - T_{800}$ should be about 10 K , but no more than 15 K
- Decided that lapse rates $> 10\text{K}/100\text{mb}$ in the lower layer are too steep, and need to be adjusted.
- Adjusted skin temperature to not exceed $10\text{K}/100\text{mb}$ limit.



Results From the Modified Computation




Bias for Alice Springs - reduced greatly; Change for Goodwin Creek - minimal




Climate Science Branch, NASA Langley Research Center



Original and Constrained Computation - Alice Springs



Bias for Alice Springs is greatly reduced




Climate Science Branch, NASA Langley Research Center




Monthly Temperature Adjustment and Frequency

Temperature Adjustment



Adjustment Frequency



Climate Science Branch, NASA Langley Research Center



Application to LW Model C

$$\text{SDLW}(\text{clr}) = a_0 + c_1 * \text{SULW} + c_2 * \ln(1 + \text{PWV}) + c_3 * [\ln(1 + \text{PWV})]^2$$

$$a_0 = 37.687, c_1 = 0.474, \\ c_2 = 94.190, c_3 = -4.935$$



$$\text{SULW} = \sigma * T_s^4$$

Use Adjusted Skin Temperature



Longwave Model A

January & July 2004 - Aqua & Terra





Climate Science Branch, NASA Langley Research Center



Longwave Model B

January & July 2004 - Aqua & Terra





Climate Science Branch, NASA Langley Research Center



Longwave Model C

January & July 2004 - Aqua & Terra




Climate Science Branch, NASA Langley Research Center



Downward Longwave Model B With MOA Ts

08 July 2004 - Terra - Daytime Overpasses





Climate Science Branch, NASA Langley Research Center



Downward Longwave Model B With Constrained Ts

08 July 2004 - Terra - Daytime Overpasses




Climate Science Branch, NASA Langley Research Center



Downward Longwave Difference Model B

08 July 2004 - Terra - (Constrained - Original)




Climate Science Branch, NASA Langley Research Center





Summary

- Determined the cause of DLF overestimation: Occurs when skin temperature (T_s) is much higher than atmospheric temperatures.
- Developed a method for detecting this condition and constraining T_s prior to using it in DLF computation. Constrained T_s is not used for upward flux computation.
- Tested the method with Model-B for all months of 2000 using an offline version.
- Applied the method to CERES versions of all three models and tested for Jan and Jul 2004 for both Terra and Aqua.
- This method remedies the overestimation. Ready to be implemented in Edition-3 β -testing and final processing.



Original and Constrained Computation - Goodwin Creek



Bias for Goodwin Creek is not affected much




Climate Science Branch, NASA Langley Research Center




Ts constrained vs Ts MOA

January & July 2004 - Aqua & Terra



• 1 • 2 - 10 • 11 - 20 • 21 - 50 • 51 - 100 • 101 - 150 • 151 - 200 • > 200




Climate Science Branch, NASA Langley Research Center



Weighting Function for DLF Reaching the Surface



(U.S. Standard Atmosphere - 50 mb Layers)



Climate Science Branch, NASA Langley Research Center



Surface and Atmospheric Temperatures Over Sea of Japan January 2000



Climate Science Branch, NASA Langley Research Center

